

BOEHRINGER INGELHEIM VETMEDICA, INC.

2621 NORTH BELT HIGHWAY, ST. JOSEPH, MO, 64506-2002

Telephone: 800-325-9167

Fax: 816-236-2717

Website: [www.bevaccinesmart.com](http://www.bevaccinesmart.com)

Website: [www.bi-vetmedica.com](http://www.bi-vetmedica.com)

Website: [www.productionvalues.us](http://www.productionvalues.us)

Website: [www.prrsresearch.com](http://www.prrsresearch.com)

Website: [www.thinkmetacam.com](http://www.thinkmetacam.com)

Website: [www.vetera-vaccines.com](http://www.vetera-vaccines.com)

Website: [www.vetmedin-us.com](http://www.vetmedin-us.com)

Website: [www.yourdogsheart.com](http://www.yourdogsheart.com)

Email: [info@productionvalues.us](mailto:info@productionvalues.us)

## BRONCHI-SHIELD® III

Boehringer Ingelheim

### **Canine Adenovirus Type 2-Parainfluenza-Bordetella Bronchiseptica Vaccine**

Modified Live Virus and Avirulent Live Culture

For vaccination of healthy dogs and puppies eight weeks of age or older as an aid in prevention of disease caused by *Bordetella bronchiseptica*, canine parainfluenza virus and canine adenovirus type 2.

**PREPARATION OF THE VACCINE:** Rehydrate with the accompanying sterile diluent. Shake well and draw back into the syringe the required amount. Remove needle from syringe and attach enclosed applicator tip. Use immediately.

**ADMINISTRATION AND DOSAGE:** Instill 0.5 mL of rehydrated vaccine into each nostril using a syringe with applicator tip. Annual revaccination is recommended.



**INFOVAX-ID® System**

The **INFOVAX-ID<sup>®</sup>** System provides a simple and effective method of recording pertinent information on the vaccines administered to animals in a veterinary practice.

For vaccines requiring reconstitution, remove label from both vials and affix both labels to the animal's medical chart.

**Using the INFOVAX-ID<sup>®</sup> System:**

U.S. Pat. No. 5,704,648

1.	Grasp the lower right hand corner of the tab at the arrow marked "Peel Here" between your thumb and forefinger ( <b>figure 1</b> ).	 <p>(fig. 1)</p>
2.	Pull steadily at a slight upward angle until the top portion of the label is separated from the vial ( <b>figure 2</b> ).	 <p>(fig. 2)</p>
3. Place the label on the animal's medical chart. Press down on the label to ensure adhesion.		

**CAUTION:** This product is designed for INTRANASAL USE ONLY. DO NOT VACCINATE DOGS PARENTERALLY. A very small percentage of animals may show sneezing, coughing or nasal discharge following vaccination. These signs are usually transient. Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. Shake well after rehydration. Burn container and all unused contents. In case of anaphylactoid reaction, administer epinephrine.

FOR INTRANASAL USE ONLY. DO NOT VACCINATE DOGS PARENTERALLY.

For Veterinary Use Only

© 2008 Fort Dodge Animal Health. All Rights Reserved.

**CAUTION:** In the absence of a veterinarian-client-patient relationship, Federal law prohibits the relabeling, repackaging, resale, or redistribution of the individual contents of this package. (9 CFR Part 112.6)

**Fort Dodge Animal Health, A division of Wyeth, Fort Dodge, Iowa 50501 USA**

U.S. Vet. License No. 112

25 DOSES	25 x 1 mL Vials of Vaccine plus 25 x 1 mL Vials of Diluent	6593 01418 1122E
----------	---	------------------------

**NAC No.:** 10281790

