

BOEHRINGER INGELHEIM VETMEDICA, INC.

2621 NORTH BELT HIGHWAY, ST. JOSEPH, MO, 64506-2002

Telephone: 800-325-9167

Fax: 816-236-2717

Website: www.bevaccinesmart.com

Website: www.bi-vetmedica.com

Website: www.productionvalues.us

Website: www.prrsresearch.com

Website: www.thinkmetacam.com

Website: www.vetera-vaccines.com

Website: www.vetmedin-us.com

Website: www.yourdogsheart.com

Email: info@productionvalues.us

DURAMUNE® MAX 5-CvK/4L THE PUPPYSHOT® BOOSTER

Boehringer Ingelheim

Canine Distemper-Adenovirus Type 2-Coronavirus-Parainfluenza-Parvovirus Vaccine

Modified Live and Killed Virus

Leptospira Canicola-Grippytyphosa-Icterohaemorrhagiae-Pomona Bacterial Extract

For subcutaneous vaccination of healthy dogs 6 weeks of age or older as an aid in the prevention of disease caused by canine distemper, infectious canine hepatitis, canine adenovirus type 2, canine coronavirus, canine parainfluenza, canine parvovirus, *Leptospira canicola*, *Leptospira icterohaemorrhagiae*, *Leptospira grippytyphosa* and *Leptospira pomona*.

DIRECTIONS FOR USE

General Directions: Aseptically rehydrate DURAMUNE Max 5 with Canine Coronavirus vaccine-Leptospira bacterial extract supplied. Administer one 1 mL dose subcutaneously.

Primary Vaccination: A recommended vaccination schedule should start at or about 6 weeks of age. The presence of maternal antibody is known to interfere with the development of active immunity. Puppies should be revaccinated every 2 to 3 weeks until 12 weeks of age.

All dogs over 12 weeks of age should initially receive one dose of Duramune Max 5-CvK/4L and a second dose 2 to 3 weeks later.

Annual Vaccination: Annual revaccination with one dose is recommended.

INFOVAX-ID[®] System

The **INFOVAX-ID[®]** System provides a simple and effective method of recording pertinent information on the vaccines administered to animals in a veterinary practice.

For vaccines requiring reconstitution, remove label from both vials and affix both labels to the animal’s medical chart.

Using the INFOVAX-ID[®] System:

U.S. Pat. No. 5,704,648

1.	Grasp the lower right hand corner of the tab at the arrow marked “Peel Here” between your thumb and forefinger (figure 1).	
2.	Pull steadily at a slight upward angle until the top portion of the label is separated from the vial (figure 2).	
3.	Place the label on the animal’s medical chart. Press down on the label to ensure adhesion.	

CAUTION: Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. In case of anaphylactoid reaction, administer epinephrine. Burn container and all unused contents.

Gentamicin and thimerosal added as preservatives.

For Veterinary Use Only

U.S. Pat. No. 5,047,238

© 2008 Fort Dodge Animal Health. All Rights Reserved.

CAUTION: In the absence of a veterinarian-client-patient relationship, Federal law prohibits the relabeling, repackaging, resale, or redistribution of the individual contents of this package. (9 CFR Part 112.6)

Fort Dodge Animal Health, A division of Wyeth, Fort Dodge, Iowa 50501 USA

U.S. Vet. License No. 112

25 DOSES	25 x 1 mL Vials of Vaccine plus 25 x 1 mL Vials of Diluent	7857
		01918
		16545D

NAC No.: 10281900