

BOEHRINGER INGELHEIM VETMEDICA, INC.

2621 NORTH BELT HIGHWAY, ST. JOSEPH, MO, 64506-2002

Telephone: 800-325-9167

Fax: 816-236-2717

Website: www.bevaccinesmart.com

Website: www.bi-vetmedica.com

Website: www.productionvalues.us

Website: www.prrsresearch.com

Website: www.thinkmetacam.com

Website: www.vetera-vaccines.com

Website: www.vetmedin-us.com

Website: www.yourdogsheart.com

Email: info@productionvalues.us

FEL-O-VAX® PCT + CALICIVAX™

Boehringer Ingelheim

Feline Rhinotracheitis-Calici-Panleukopenia Vaccine

Killed Virus

Includes hemorrhagic calici strain

For subcutaneous vaccination of healthy cats 8 to 10 weeks of age or older, as an aid in the prevention of disease caused by feline rhinotracheitis, calici and panleukopenia viruses. The vaccine contains one additional feline calicivirus strain isolated from one hemorrhagic calicivirus outbreak that occurred in 1998 and was shown to protect vaccinated cats against calicivirus-induced hemorrhagic disease in controlled vaccination challenge experiments.

DOSE: Inject one 1 mL dose subcutaneously using aseptic technique. Repeat in 3 to 4 weeks. Annual revaccination with a single dose of vaccine is recommended.



INFOVAX-ID® System

The **INFOVAX-ID[®]** System provides a simple and effective method of recording pertinent information on the vaccines administered to animals in a veterinary practice.

For vaccines requiring reconstitution, remove label from both vials and affix both labels to the animal's medical chart.

Using the INFOVAX-ID[®] System:

U.S. Pat. No. 5,704,648

| | |
|----------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------|
| <p>1. Grasp the lower right hand corner of the tab at the arrow marked "Peel Here" between your thumb and forefinger (figure 1).</p> |  <p>(fig. 1)</p> |
| <p>2. Pull steadily at a slight upward angle until the top portion of the label is separated from the vial (figure 2).</p> |  <p>(fig. 2)</p> |
| <p>3. Place the label on the animal's medical chart. Press down on the label to ensure adhesion.</p> | |

CAUTION: Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. Use entire contents when first opened (10 mL). In case of anaphylactoid reaction, administer epinephrine.

Thimerosal, neomycin, polymyxin B and amphotericin B added as preservatives.

For Veterinary Use Only

U.S. Patent Nos. 5,047,238 and patents pending.

© 2006, 2009 Fort Dodge Animal Health. All Rights Reserved.

CAUTION: In the absence of a veterinarian-client-patient relationship, Federal law prohibits the relabeling, repackaging, resale, or redistribution of the individual contents of this package. (9 CFR Part 112.6)

Fort Dodge Animal Health, A division of Wyeth, Fort Dodge, Iowa 50501 USA

U.S. Vet. License No. 112

| | | |
|----------|----------------------------|--------|
| 50 DOSES | 50 - 1 mL Vials of Vaccine | 18366B |
|----------|----------------------------|--------|

NAC No.: 10282570