

VAN BEEK NATURAL SCIENCE, LLC.

3689 460TH STREET, ORANGE CITY, IA, 51041

Telephone: 712-737-2958

Toll-Free: 800-346-5311

Fax: 712-737-2878

Website: www.vanbeeknaturalscience.com

Email: info@vanbeeknaturalscience.com

DIAGEL™ EQUINE

Van Beek

Diarrhea Control Gel

Equine (Horse / Foal)

National Animal Supplement Council - THE BENCHMARK OF QUALITY™

Product Facts:

Active Ingredients per ml:

Proprietary Blend of (Carvacrol, Cinnameldahyde, Eugenol, Thymol)	51 mg
Sesquiterpenoid	3.5 mg

Inactive Ingredients:

Anise Oil, Arabic Gum, Carrageenan, Cellulose Gum, Deionized Water, Lecithin, Polysorbate 80, Potassium Chloride, Propylene Glycol, Sodium Chloride, Sodium Hexametaphosphate.

Cautions:

Safe use in pregnant animals or animals intended for breeding has not been proven. If animal's condition worsens or does not improve, stop product administration and consult your veterinarian.

Veterinary Use Only

FOR USE IN EQUINE ONLY

Recommended for support in controlling environmentally induced diarrhea.

Primary Use: *DiaGel* is an orally administered product which may aid in controlling bacterial diarrhea. *DiaGel* eliminates bacteria such as *E. coli* and *Salmonella* which occur naturally in the GI tract.

Causes of bacterial diarrhea may include:

- Foal Stress
- Dietary changes (grain overload)
- Environmental Changes
- Secondary Stress

A one-time administration may firm stools within 24 hours.

Secondary Use: *DiaGel* may be used in conjunction with fluid therapy, endotoxin therapy, or antibiotics to treat secondary infections associated with other causes of diarrhea.

Pre-Stress Use: *DiaGel* may be administered 2 hours prior to stress-induced events (racing, showing, trailering, stabling, or foaling) to help reduce the incidence of diarrhea.

Directions for use:

Set the calibration ring on the syringe plunger according to the weight usage requirement. To administer, place the syringe between the side teeth of the horse, aiming as far down on the back of the tongue as possible. Gently push the plunger to release the gel.

Usage: Referencing the appropriate use, administer *DiaGel* according to the horse's weight:

10 ml - Newborn to 120 pounds

15 ml - 121 to 250 pounds

20 ml - 251 to 500 pounds

30 ml - 501+ pounds

Primary Use: Administer the recommended amount by weight for bacterial diarrhea in the GI tract. Most horses respond in 12 hours with the majority of horses responding in 24 hours or less. If necessary, double the usage for the initial administration. If additional administrations are necessary, use the recommended amount by weight every 6 hours for up to 3 days.

Secondary Use: Administer the recommended amount by weight every 12 hours for up to 5 days.

Pre-Stress Use: Administer the recommended amount by weight 2 hours prior to stress induced event. Repeat as necessary by weight every 12 hours for up to 5 days.

Warnings:

For animal use only. Keep out of the reach of children and other animals. In case of accidental overdose, contact a health professional immediately. This product should not be given to animals intended for human consumption.

Storage:

Store in a cool, dry place protected from light.

Questions?

Van Beek Natural Science, Orange City, IA

1-800-346-5311

www.vanbeeknaturalscience.com

US Patent No. 6,414,036

Made in USA

Net contents:		
1 (30ml) syringe	TY#PBG021	111708

NAC No.: 61620031