

Fish Cin (Clindamycin 150mg), 60 Tablets

Clindamycin Antibiotics for Fish

Fish Cin is for use in aquariums and exerts a bactericidal action on gram positive and some gram negative bacteria. Useful for control of some common bacterial fish diseases.

Off-Label Use in Dogs & Cats: Metronidazole is a antibacterial, antiprotozoal agent that has been used in the treatment of giardiasis, trichomoniasis, amoebiasis, balantidiasis, and trypanosomiasis. It is as effective and is better tolerated than quinacrine for the treatment of giardiasis. It is bactericidal to many anaerobic bacteria. It has been used in the treatment of septicemia, meningitis, peritonitis, biliary infections, colitis, and gingivostomatitis in which anaerobic bacteria are involved. The drug may also have immunosuppressive or immunostimulatory properties.

Directions:

Fish: Add one tablet (150mg) into aquarium for each 10 gallons of water. Repeat in 24 hours. It is suggested that a partial water change be made between treatments. While treatment duration depends on the type and severity of infection, it is recommended that extended baths in the medication continue for a minimum of 5 days and for not more than 10 days. Discontinued treatment if no improvement is noted within 5 days.

Dogs and Cats: Fish Cin is commonly given in off-label use to small animals.

Anaerobic Infection

Dog: 11 mg/Kg (5 mg/lb) every 12 hours orally.

Cat: 11 mg/Kg (5 mg/lb) every 12 hours orally.

Staphylococcal Infections

Dog: 11 mg/Kg (5 mg/lb) every 12 hours, or 22 mg/Kg (10 mg/lb) every 24 hours.

Cat: 5.5 mg/Kg (2.5mg/lb) every 12 hours, or 11 mg/Kg (5 mg/lb) every 24 hours.

Toxoplasmosis

Dog: 12.5 mg/Kg (5.7 mg/lb) every 12 hours for 4 weeks.

Cat: 12.5 mg/Kg (5.7 mg/lb) every 12 hours for 4 weeks.

Do not administer to rabbits or rodents because it causes diarrhea. Do not administer to horses or ruminants because it could cause severe enteritis.

Not for human use.