

BUTLER SCHEIN™ ANIMAL HEALTH

A Henry Schein Company

400 METRO PLACE NORTH, DUBLIN, OH, 43017-7545

Telephone: 614-761-9095

Toll-Free: 1-888-691-2724 (My1-BSAH)

Website: www.ButlerSchein.com

PARID EQ™ PASTE 1.87%

Butler Schein™ Animal Health

(ivermectin)

Anthelmintic And Boticide

Removes worms and bots with a single dose.

Contents will treat up to 1250 lb body weight

For Oral Use In Horses Only

ANADA 200-286, Approved by FDA

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Parid EQ™ (ivermectin) Paste 1.87% provides effective control of the following parasites in horses: **Large Strongyles** (adults) - *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp; **Small Strongyles** including those resistant to some benzimidazole class compounds (adults and fourth-stage larvae) - *Cyathostomum* spp, *Cylicocyclus* spp; *Cylicostephanus* spp; *Cylicodontophorus* spp; **Pinworms** (adults and fourth-stage larvae) - *Oxyuris equi*; **Ascarids** (adults and third- and fourth-stage larvae) - *Parascaris equorum*; **Hairworms** (adults) - *Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults) - *Habronema muscae*; **Bots** (oral and gastric stages) - *Gastrophilus* spp; **Lungworms** (adults and fourth-stage larvae) - *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults) - *Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp cutaneous third-stage larvae; **Dermatitis** caused by neck threadworm microfilariae, *Onchocerca* sp.

DOSAGE AND ADMINISTRATION: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight. (1) While holding plunger, turn the knurled ring on the plunger to the right so the side nearest the barrel is at the prescribed weight marking. (2) Make sure that the horse's mouth contains no feed. (3) Remove the cover from the tip of the syringe. (4) Insert the syringe

tip into the horse's mouth at the space between the teeth. (5) Depress the plunger as far as it will go, depositing paste on the back of the tongue. (6) Immediately raise the horse's head for a few seconds after dosing.

PARASITE CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate.

Consult your veterinarian for a control program to meet your specific needs. Parid EQ™ (ivermectin) Paste 1.87% effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *S. vulgaris*.

PRODUCT ADVANTAGES: Broad -Spectrum Control-Parid EQ™ (ivermectin) Paste 1.87% kills important internal parasites, including bots and the arterial stages of *Strongylus vulgaris*, with a single dose. Parid EQ™ (ivermectin) Paste 1.87% is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

Safety-Parid EQ™ (ivermectin) Paste 1.87% may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

▶ **RESIDUE WARNING:** Do not use in horses intended for food purposes. ◀

CAUTION: Parid EQ™ (ivermectin) Paste 1.87% has been formulated specifically for use in horses **only**. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result. Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. **Keep this and all drugs out of the reach of children.** Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

NOTE TO USER: Swelling and itching reactions after treatment with Parid EQ™ (ivermectin) Paste 1.87% have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp) microfilariae. These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with Parid EQ™ (ivermectin) Paste 1.87%. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

EACH SYRINGE CONTAINS 6.08 g (0.21 oz) IVERMECTIN PASTE

Store at controlled room temperature, 20° to 25°C (68° to 77°F).

800003

Made in U.S.A.

Manufactured by IVX Animal Health, Inc., St. Joseph, MO 64503

Manufactured for: BUTLER ANIMAL HEALTH SUPPLY, DUBLIN, OHIO 43017

Net Wt.	
---------	--

6.08 g (0.21 oz)	Iss. 01-03
------------------	------------

NAC No.: 10822083