

BOEHRINGER INGELHEIM VETMEDICA, INC.

2621 NORTH BELT HIGHWAY, ST. JOSEPH, MO, 64506-2002

Telephone: 800-325-9167

Fax: 816-236-2717

Website: [www.bevaccinesmart.com](http://www.bevaccinesmart.com)

Website: [www.bi-vetmedica.com](http://www.bi-vetmedica.com)

Website: [www.productionvalues.us](http://www.productionvalues.us)

Website: [www.prrsresearch.com](http://www.prrsresearch.com)

Website: [www.thinkmetacam.com](http://www.thinkmetacam.com)

Website: [www.vetera-vaccines.com](http://www.vetera-vaccines.com)

Website: [www.vetmedin-us.com](http://www.vetmedin-us.com)

Website: [www.yourdogsheart.com](http://www.yourdogsheart.com)

Email: [info@productionvalues.us](mailto:info@productionvalues.us)

## **DURAMUNE LYME® + MAX 5-CvK**

Boehringer Ingelheim

### **Canine Distemper-Adenovirus Type 2-Coronavirus-Parainfluenza-Parvovirus Vaccine**

Modified Live and Killed Virus

### **Borrelia Burgdorferi Bacterin**

For Veterinary Use Only

For subcutaneous vaccination of healthy dogs 9 weeks of age or older as an aid in the prevention of disease caused by canine distemper, infectious canine hepatitis, canine adenovirus type 2, canine coronavirus, canine parainfluenza, canine parvovirus and *Borrelia burgdorferi*.

### **Directions for Use**

**- General Directions:** Aseptically rehydrate Duramune Max 5 with Canine Coronavirus vaccine, *Borrelia burgdorferi* bacterin supplied. Administer (one) 1 mL dose subcutaneously.

The presence of maternal antibody is known to interfere with the development of active immunity. Puppies should be revaccinated every 2 to 3 weeks until 12 weeks of age.



All dogs over 12 weeks of age should initially receive one dose of this product and a second dose 2 to 3 weeks later.

**- Annual Vaccination:** Annual revaccination with one dose is recommended.

The **PEEL-OFF LABEL I.D. SYSTEM** provides a simple and effective method of recording pertinent information on the vaccines administered to animals in a veterinary practice.

For vaccines requiring reconstitution, remove label from both vials and affix both labels to the animal's medical chart.

**Using the System:**

<p>1. Grasp the vial label at the corner marked with an indicator arrow, between your thumb and forefinger (<b>figure 1</b>).</p>	 <p>(fig. 1)</p>
<p>2. Pull steadily at a slight angle until the label is separated from the vial (<b>figure 2</b>).</p>	 <p>(fig. 2)</p>
<p>3. Place the label on the animal's medical chart. Press down on the label to ensure adhesion.</p>	

**Caution:** Store in the dark at 2° to 7°C (35° to 45°F). AVOID FREEZING. SHAKE WELL. In case of anaphylactoid reaction, administer epinephrine. Burn container and all unused contents.

Gentamicin and thimerosal added as preservatives.

**Caution:** In the absence of a veterinarian-client-patient relationship, Federal law prohibits the relabeling, repackaging, resale, or redistribution of the individual contents of this package.

Duramune Lyme® is a registered trademark of Boehringer Ingelheim Vetmedica, Inc.

© 2011 Boehringer Ingelheim Vetmedica, Inc. All Rights Reserved.

**Boehringer Ingelheim Vetmedica, Inc., St. Joseph, MO 64506 U.S.A.**

**U.S. Veterinary License No. 124**

25 Doses	25 Vials of Vaccine plus 25 x 1 mL Vials of Diluent	D3381B/D9161B
----------	---	---------------

**NAC No.:** 10282501

