

Preventic Tick Collar for Dogs

Product Description:

Kills and detaches ticks before they can transfer disease-causing organisms. Also recommended for the treatment of Demodex in dogs.

Indications:

Tick collar for dogs. Kills ticks for 3 months.

Directions for Use:

It is a violation of Federal Law to use this collar in a manner inconsistent with its labeling.

Read entire label directions before each use.

Place the PREVENTIC® collar around the dog's neck, buckle and adjust for proper fit. Cut off and dispose of excess length. Do not allow dogs to play with cut-off excess. The collar must be worn tightly enough to contact skin and so that the dog cannot remove it. Activity against ticks starts within 24 hours and continues as long as the collar is worn (up to 90 days). Ticks present in the dog's environment that may reappear on the dog will not attach and will be killed in less than 24 hours. Replace collar when effectiveness diminishes. The effectiveness of the collar is not diminished by normal wetting such as rainfall, however, it is suggested that the collar be removed before bathing. The active ingredient is not an insecticide. Therefore, use other means for the control of fleas.

Active Ingredient(s):

Amitraz: N'-(2,4-dimethylphenyl)-N-[[2,4-dimethylphenyl)-imino]methyl]-N-methylmethanimidamide	9.0%
Inert Ingredients	91.0%

Contraindication(s):

Use only on dogs. Do not use on cats.

Precautionary Statements:

Hazard to Humans and Domestic Animals:

Caution: This collar is intended for use as an acaricide generator and is not to be taken internally by man or animal. Not intended for use on humans. Do not allow children to handle or play with this collar.

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Some humans may develop skin irritation through contact with collared pet. Wash thoroughly with soap and water after handling.

Do not use on puppies less than 12 weeks old. Consult a veterinarian before using on debilitated, aged, medicated, pregnant or nursing animals. Do not treat collared dogs with other monoamine oxidase inhibitors, or with tricyclic antidepressants, selective serotonin reuptake inhibitors or pressor agents. Some dogs may be sensitive to this collar. Remove at first sign of irritation or adverse reaction, bathe dog with a non-pesticidal shampoo and rinse with large amounts of water. If signs continue, consult a veterinarian immediately. There is a risk of collar breakage and ingestion if collar is worn by unattended dogs that play by mouthing each other in the neck area. If collar is missing from a dog, search for, retrieve and recover the entire collar. Immediately consult a veterinarian if dog is depressed.

Note to Veterinarian: If a dog ingests a PREVENTIC® collar and exhibits intoxication (primary sign is severe depression), induce vomiting, administer non-oily laxative, activated charcoal and enema. Provide supportive therapy until collar fragments are voided. If dog cannot be roused, Yohimbine, 0.1 mg/kg by intravenous injection is antidotal. Repeat as needed.

Disposal:

Do not reuse empty pouch, box or collar. Wrap in newspaper and put in trash collection.

Warning:

Keep out of reach of children.